

First Passage Percolation Has Sublinear Distance Variance

Itai Benjamini Gil Kalai Oded Schramm

October 21, 2002

Abstract

Let $0 < a < b < \infty$, and for each edge e of $\mathbb{Z}^d$ let $\omega_e = a$ or $\omega_e = b$, each with probability $1/2$, independently. This induces a random metric dist_ω on the vertices of $\mathbb{Z}^d$, called first passage percolation. We prove that for $d > 1$ the distance $\text{dist}_\omega(0, v)$ from the origin to a vertex v , $|v| > 2$, has variance bounded by $C |v| / \log |v|$, where $C = C(a, b, d)$ is a constant which may only depend on a , b and d . Some related variants are also discussed.

1 Introduction

Consider the following model of first passage percolation. Fix some $d = 2, 3, \dots$, and let $E = E(\mathbb{Z}^d)$ denote the set of edges in $\mathbb{Z}^d$. Also fix numbers $0 < a < b < \infty$. Let $\Omega := \{a, b\}^E$ carry the product measure, where $\mathbf{P}[\omega_e = a] = \mathbf{P}[\omega_e = b] = 1/2$ for each $e \in E$. Given $\omega = (\omega_e : e \in E) \in \Omega$ and vertices $v, u \in \mathbb{Z}^d$, let $\text{dist}_\omega(v, u)$ denote the least distance from v to u in the metric induced by ω ; that is, the infimum of $\text{length}_\omega(\alpha) := \sum_{e \in \alpha} \omega_e$, where α ranges over all finite paths in $\mathbb{Z}^d$ from u to v . Let $|v| := \|v\|_1$ for vertices $v \in \mathbb{Z}^d$.

Theorem 1. *There is a constant $C = C(d, a, b)$ such that for every $v \in \mathbb{Z}^d$, $|v| \geq 2$,*

$$\text{var}(\text{dist}_\omega(0, v)) \leq C \frac{|v|}{\log |v|}.$$

In [12] Kesten used martingale inequalities to prove $\text{var}(\text{dist}_\omega(0, v)) \leq C|v|$ and proved some tail estimates. Talagrand [17] used his “convexified” discrete isoperimetric inequality to prove that for all $t > 0$,

$$\mathbf{P}\left[|\text{dist}_\omega(0, v) - M| \geq t\sqrt{|v|}\right] \leq C \exp(-t^2/C), \quad (1)$$

where M is the median value of $\text{dist}_\omega(0, v)$. (Both Kesten’s and Talagrand’s results apply to more general distributions of the edge lengths ω_e .) “Novice readers might expect to hear next of a central limit theorem being proved,” writes Durrett [7], describing Kesten’s results, “however physicists tell us ... that in two dimensions the standard deviation ... is of order $|v|^{1/3}$.” Recent remarkable work [1, 9, 6] not only supports this prediction, but suggests what the limiting distribution and large deviation behavior is. The case of a certain variant of **oriented** first passage percolation is settled by Johansson [9]. For lower bounds on the variance in two-dimensional first passage percolation see Newman-Piza [14] and Pemantle-Peres [15].

As in Kesten’s and Talagrand’s earlier results, the most essential feature about first passage percolation which we use is that the number of edges $e \in E$ such that modifying ω_e increases $\text{dist}_\omega(0, v)$ is bounded by $C|v|$.

Another essential ingredient in the current paper is the following extension by Talagrand [16] of an inequality by Kahn, Kalai and Linial [10]. Let J be a finite index set. For $j \in J$, and $\omega \in \{a, b\}^J$ let $\sigma_j \omega$ be the element of $\{a, b\}^J$ which is different from ω only in the j -th coordinate. For $f : \{a, b\}^J \rightarrow \mathbb{R}$ set $\sigma_j f := f \circ \sigma_j$ and

$$\rho_j f(\omega) := \frac{f(\omega) - \sigma_j f(\omega)}{2}.$$

Talagrand’s [16, Thm. 1.5] inequality is

$$\text{var}(f) \leq C \sum_{j \in J} \frac{\|\rho_j f\|_2^2}{1 + \log(\|\rho_j f\|_2 / \|\rho_j f\|_1)}, \quad (2)$$

where C is a universal constant. (In Section 4 we supply a direct proof of (2) from the Bonami-Beckner inequality. A very reasonable upper bound for C can be obtained from this proof. We also explain there how (2) easily implies (1).)

The basic idea in the proof of Theorem 1 is to apply this inequality to $f(\omega) := \text{dist}_\omega(0, v)$. For this, we wanted to show that, roughly, $\mathbf{P}[\rho_e f(\omega) \neq 0]$

is small, except for a small number of edges e . However, since we were not able to prove this, we had to resort to an averaging trick.

Theorem 1 should hold for other models of first passage percolation, where the edge lengths have more general distributions. The relevant result of [10], as well as (2) that we use rely on the Bonami-Beckner [4, 2] inequality, which holds for $\{a, b\}^n$, but fails on some more general product spaces. The paper [5] does extend some of these results to general product spaces; see also [8]. Talagrand's [16, Thm. 1.5] applies to product measures on $\{a, b\}^n$, which are not necessarily uniform. Beckner [2] uses his inequality to derive a similar result for the Gaussian measure on $\mathbb{R}^n$, and an analog for Talagrand's inequality (2) for the Gaussian measure (pointed out to us by Assaf Naor) can be found in [3]. In the present paper, we preferred simplicity to generality, but it will be interesting to extend the theorem to more general distributions. We refer the reader also to the book by Ledoux [13] for a general view of relevant techniques and knowledge.

It seems even more important to put effort into the fundamental task of lowering the upper bound from $C|v|/\log|v|$ to $C|v|^{1-\epsilon}$, $\epsilon > 0$.

To first illustrate the basic technique in a simpler setting, we will also prove a theorem about the variance of the first passage percolation circumference of a discrete torus, or more generally in a cartesian product of a finite vertex-transitive graph with a circle.

Let H be a finite vertex-transitive graph, and let $n \in \mathbb{N}_+$. Let G be the product of H and the cycle of length n , $\mathbb{Z}/(n\mathbb{Z})$. We say that a closed path γ in G is a circumference, if its projection onto $\mathbb{Z}/(n\mathbb{Z})$ has degree 1; namely, γ can be oriented so that its projection has one more edge going from 0 to 1 than from 1 to 0. If $\omega : E(G) \rightarrow \{a, b\}$, define its circumference length $c_G(\omega)$ as the minimal ω -length of any circumference path.

Theorem 2. *Let $n \in \mathbb{N}_+$ and let G be a cartesian product of a vertex-transitive graph H with the cycle $\mathbb{Z}/(n\mathbb{Z})$ of length n . Let $\omega \in \{a, b\}^{E(G)}$ be random-uniform. Then*

$$\text{var}(c_G(\omega)) \leq C \frac{b}{a} (b-a)^2 \frac{n}{1 + \log(a|V(H)|/b)},$$

where C is a universal constant.

For example, when G is the square torus $G = (\mathbb{Z}/(n\mathbb{Z}))^2$, with $n > 1$, we get the estimate $\text{var}(c_G) \leq C (b/a) (b-a)^2 n / \log n$.

2 Proof of Theorem 2

Let β be a circumference path in G such that $\sum_{e \in \beta} \omega_e = c_G(\omega)$. We use some arbitrary but fixed method for choosing between all possible choices for β . Clearly, $c_G(\omega) \leq bn$. Therefore,

$$|\beta| \leq bn/a, \quad (3)$$

where $|\beta|$ denotes the number of edges in β .

Let $e \in E(G)$. Note that if $\rho_e c_G(\omega) < 0$, then we must have $e \in \beta$. By (3), this gives

$$\sum_{e \in E(G)} \mathbf{P}[\rho_e c_G < 0] \leq \mathbf{E}[|\beta|] \leq bn/a. \quad (4)$$

Let Γ be the automorphism group of G . Fix $e \in E(G)$. By symmetry, $\mathbf{P}[\rho_e c_G < 0] = \mathbf{P}[\rho_{e'} c_G < 0]$ for every $e' \in \Gamma e$. Consequently,

$$|\Gamma e| \mathbf{P}[\rho_e c_G < 0] \leq \sum_{e' \in E(G)} \mathbf{P}[\rho_{e'} c_G < 0] \leq bn/a.$$

Now note that since H is vertex-transitive, also G is vertex-transitive, and consequently $|\Gamma e| \geq |V(G)|/2 = n|V(H)|/2$. Therefore,

$$\mathbf{P}[\rho_e c_G < 0] \leq b|V(H)|^{-1}/a. \quad (5)$$

Now clearly, $\mathbf{P}[\rho_e c_G \neq 0] = 2\mathbf{P}[\rho_e c_G < 0]$ and $\|\rho_e c_G\|_\infty \leq (b-a)/2$. Therefore,

$$\|\rho_e c_G\|_2^2 \leq (1/2)(b-a)^2 \mathbf{P}[\rho_e c_G < 0]. \quad (6)$$

By Cauchy-Schwarz,

$$\|\rho_e c_G\|_1 \leq \sqrt{\mathbf{P}[\rho_e c_G \neq 0]} \|\rho_e c_G\|_2. \quad (7)$$

By (2), we have

$$\text{var}(c_G) \leq C \frac{\sum_{e \in E(G)} \|\rho_e c_G\|_2^2}{1 + \min_{e \in E(G)} \log(\|\rho_e c_G\|_2 / \|\rho_e c_G\|_1)}.$$

To estimate the numerator, we use (6) and (4), and for the denominator (5) and (7). The theorem easily follows. $\square$

3 Proof of Theorem 1

If $v, u \in \mathbb{Z}^d$, and α is a path from v to u , then α will be called an ω -**geodesic** if it minimizes ω -length; that is, $\text{dist}_\omega = \sum_{e \in \alpha} \omega_e$. Given ω , let γ be an ω -geodesic from 0 to v . Although there may be more than one such geodesic, we require that γ depends only on ω . (For example, we may use an arbitrary deterministic choice among any possible collection of ω -geodesics.)

The general strategy for the proof of Theorem 1 is as for Theorem ???. However, the difficulty is that there is not enough symmetry to get a good bound on $\mathbf{P}[\rho_e \text{dist}_\omega(0, v) < 0]$. It would have been enough to show that $\mathbf{P}[e \in \gamma] < C |v|^{-1/C}$ holds with the exception of at most $C |v| / \log |v|$ edges, for some constant $C > 0$. But we could not prove this. Therefore, we will need an averaging argument, for which the following lemma will be useful.

Lemma 3. *There is a constant $c > 0$ such that for every $m \in \mathbb{N}$ there is a function*

$$g = g_m : \{a, b\}^{m^2} \rightarrow \{0, 1, 2, \dots, m\}$$

satisfying

$$\|\sigma_j g - g\|_\infty \leq 1$$

for every $j = 1, 2, \dots, m^2$ and

$$\max_y \mathbf{P}[g(x) = y] \leq c/m,$$

where x is random-uniform in $\{a, b\}^{m^2}$.

PROOF: Assume $a = 0$ and $b = 1$, for simplicity of notation. Let $k : \mathbb{N} \rightarrow \{0, 1, 2, \dots, m\}$ be the function satisfying $k(0) = 0$, $k(j+1) = k(j) + 1$ when $j \in \bigcup_{s=0}^\infty [2sm, 2sm + m - 1]$ and $k(j+1) = k(j) - 1$ for all other $j \in \mathbb{N}$. It is left to the reader to check that $g(x) = k(\|x\|_1)$ has the required properties. $\square$

Fix some $v \in \mathbb{Z}^d$ with $|v|$ large, and set $f = f(\omega) := \text{dist}_\omega(0, v)$. Clearly, $f(\omega) \leq b|v|$, and therefore $|\gamma| \leq (b/a)|v|$, where $|\gamma|$ denotes the number of edges in γ . In particular, f depends on only finitely many of the coordinates in ω . Also, $|\gamma| \leq (b/a)|v|$ implies

$$\sum_{e \in E} \mathbf{P}[e \in \gamma] = \mathbf{E}[|\gamma|] \leq (b/a)|v|. \quad (8)$$

Fix $m := \lfloor |v|^{1/4} \rfloor$, and let $S := \{1, \dots, d\} \times \{1, \dots, m^2\}$. Let $c > 0$ and $g = g_m$ be as in Lemma 3. Given $x = (x_{i,j} : \{i, j\} \in S) \in \{0, 1\}^S$ let

$$z = z(x) := \sum_{i=1}^d g(x_{i,1}, \dots, x_{i,m^2}) \mathbf{e}_i,$$

where $\{\mathbf{e}_1, \dots, \mathbf{e}_d\}$ is the standard basis for $\mathbb{R}^d$. Define

$$\tilde{f}(x, \omega) := \text{dist}_\omega(z, v + z).$$

We think of $\tilde{f}$ as a function on the space $\tilde{\Omega} := \{a, b\}^{S \cup E}$. Since $|z| \leq md$, it follows that $|\tilde{f} - f| \leq 2mdb$. In particular,

$$\text{var}(f) \leq \text{var}(\tilde{f}) + 4mdb\sqrt{\text{var}(\tilde{f})} + 4m^2d^2b^2. \quad (9)$$

It therefore suffices to find a good estimate for $\text{var}(\tilde{f})$.

Let $e \in E$ be some edge. We want to estimate its influence:

$$I_e(\tilde{f}) := \mathbf{P}[\sigma_e \tilde{f}(x, \omega) \neq \tilde{f}(x, \omega)] = 2\mathbf{P}[\sigma_e \tilde{f}(x, \omega) > \tilde{f}(x, \omega)].$$

(Here, $\mathbf{P}$ is the uniform measure on $\tilde{\Omega}$.) Note that if the pair $(x, \omega) \in \tilde{\Omega}$ satisfies $\sigma_e \tilde{f}(x, \omega) > \tilde{f}(x, \omega)$, then e must be on every ω -geodesic from z to $v + z$. Consequently, conditioning on z and translating ω and e by $-z$ gives

$$I_e(\tilde{f}) = 2\mathbf{P}[\sigma_e \tilde{f}(x, \omega) > \tilde{f}(x, \omega)] \leq 2\mathbf{P}[e - z \in \gamma]. \quad (10)$$

Let Q be the set of edges $e' \in E(\mathbb{Z}^d)$ such that $\mathbf{P}[e - z = e'] > 0$. The L^1 diameter of Q is $O(m)$. (We allow the constants in the $O(\cdot)$ notation to depend on d, a and b , but not on v .) Hence, the diameter of Q in the dist_ω metric is also $O(m)$, and therefore $|\gamma \cap Q| \leq O(m)$. But the lemma gives

$$\max_{z_0} \mathbf{P}[z = z_0 \mid \omega] \leq (c/m)^d.$$

By conditioning on γ and summing over the edges in $\gamma \cap Q$, we therefore get

$$\mathbf{P}[e \in \gamma + z \mid \gamma] \leq O(1) m^{1-d}.$$

Consequently, (10) and the choice of m give

$$I_e(\tilde{f}) \leq O(1) |v|^{-1/4}. \quad (11)$$

Also, (8) implies

$$\sum_{e \in E} \mathbf{P}[e - z \in \gamma \mid z] \leq (b/a) |v|.$$

Combining this with (10) therefore gives

$$\sum_{e \in E} I_e(\tilde{f}) \leq 2(b/a) |v|.$$

Applying (11) yields

$$\sum_{e \in E} \frac{I_e(\tilde{f})}{1 + |\log I_e(\tilde{f})|} \leq O(1) |v| / \log |v|. \quad (12)$$

On the other hand, $I_s(\tilde{f}) \leq 2(b-a)$ for $s \in S$. As $|S| = O(1) |v|^{1/2}$ and $\|\rho_q f\|_\infty = O(1)$ for $q \in S \cup E$, we get from (12) and (2)

$$\text{var}(\tilde{f}) \leq O(1) \sum_{q \in E \cup S} \frac{I_q(\tilde{f})}{1 + |\log I_q(\tilde{f})|} \leq O(1) |v| / \log |v|.$$

Therefore, Theorem 1 now follows from (9). $\square$

As alluded to in the introduction, the proof would have been simpler if we could show that there is a $C > 0$ such that $\mathbf{P}[\rho_e f(\omega) \neq 0] < |v|^{-C}$ holds with the possible exception of $|v| / \log |v|$ edges. It would be interesting to prove the closely related statement that the probability that γ passes within distance 1 from $v/2$ tends to zero as $|v| \rightarrow \infty$.

4 A proof of Talagrand's inequality (2)

To prove (2), it clearly suffices to take $a = 0, b = 1$. For $f : \{0, 1\}^J \rightarrow \mathbb{R}$, consider the Fourier-Walsh expansion of f ,

$$f = \sum_{S \subset J} \widehat{f}(S) u_S,$$

where $u_S(\omega) = (-1)^{S \cdot \omega}$ and $S \cdot \omega$ is shorthand for $\sum_{s \in S} \omega_s$. For each $p \in \mathbb{R}$ define the operator

$$T_p(f) := \sum_{S \subset J} p^{|S|} \widehat{f}(S) u_S,$$

which is of central importance in harmonic analysis. The Bonami-Beckner [4, 2] inequality asserts that

$$\|T_p f\|_2 \leq \|f\|_{1+p^2}. \quad (13)$$

Set $f_j := \rho_j f$. Because $\rho_j u_S = u_S$ if $j \in S$ and $\rho_j u_S = 0$ if $j \notin S$, we have

$$\widehat{f_j}(S) = \begin{cases} \widehat{f}(S) & j \in S, \\ 0 & j \notin S. \end{cases}$$

Since $\|g\|_2^2 = \sum_{S \subset J} \widehat{g}(S)^2$, it follows that

$$\int_0^1 \|T_p f_j\|_2^2 dp = \int_0^1 \sum_{S \subset J} p^{2|S|} \widehat{f_j}(S)^2 dp = \sum_{S \subset J} 1_{j \in S} \frac{\widehat{f}(S)^2}{2|S|+1}.$$

Consequently,

$$\sum_{j \in J} \int_0^1 \|T_p f_j\|_2^2 dp = \sum_{S \subset J} |S| \frac{\widehat{f}(S)^2}{2|S|+1} \geq \frac{1}{3} \sum_{\emptyset \neq S \subset J} \widehat{f}(S)^2 = \frac{\text{var}(f)}{3}.$$

Therefore, (13) gives

$$\text{var}(f) \leq 3 \sum_{j \in J} \int_0^1 \|f_j\|_{1+p^2}^2 dp. \quad (14)$$

An instance of the Hölder inequality

$$\mathbf{E}[|f_j|^{1+p^2}] \leq \mathbf{E}[f_j^2]^{p^2} \mathbf{E}[|f_j|]^{1-p^2}$$

implies

$$\begin{aligned} \int_0^1 \|f_j\|_{1+p^2}^2 dp &\leq \int_0^1 \left(\mathbf{E}[f_j^2]^{p^2} \mathbf{E}[|f_j|]^{1-p^2} \right)^{2/(1+p^2)} dp \\ &= \|f_j\|_2^2 \int_0^1 (\|f_j\|_1 / \|f_j\|_2)^{2(1-p^2)/(1+p^2)} dp \\ &\leq 2 \|f_j\|_2^2 \int_{1/2}^1 (\|f_j\|_1 / \|f_j\|_2)^{2(1-p^2)/(1+p^2)} dp. \end{aligned}$$

Let $s(p) := 2(1 - p^2)/(1 + p^2)$. Since $s'(p) \leq s'(1) = -2$ when $p \in [1/2, 1]$, the above gives

$$\begin{aligned} \int_0^1 \|f_j\|_{1+p^2}^2 dp &\leq 2 \|f_j\|_2^2 \int_{s(1/2)}^{s(1)} (\|f_j\|_1 / \|f_j\|_2)^s \frac{ds}{s'(p)} \\ &\leq \|f_j\|_2^2 \int_0^{6/5} (\|f_j\|_1 / \|f_j\|_2)^s ds \\ &= \|f_j\|_2^2 \frac{1 - (\|f_j\|_1 / \|f_j\|_2)^{6/5}}{\log(\|f_j\|_2 / \|f_j\|_1)}. \end{aligned}$$

Now (14) implies

$$\text{var}(f) \leq 3 \sum_{j \in J} \|f_j\|_2^2 \frac{1 - (\|f_j\|_1 / \|f_j\|_2)^{6/5}}{\log(\|f_j\|_2 / \|f_j\|_1)} \quad (15)$$

from which (2) follows. $\square$

Remark: It is worth noting that the tail estimate (1) can be derived from the variance inequality (2). Indeed, let $f = \text{dist}_\omega(0, v)$ and $u \in (0, 3/4)$. Set $s = s(u) := \inf\{t \geq 0 : \mathbf{P}[f(\omega) > t] < u\}$, and define $g(\omega) := \max\{f(\omega), s\}$. It follows from (2) that

$$\text{var}(g) \leq C u |v| / (1 + \min_e \log(\|\rho_e g\|_2 / \|\rho_e g\|_1)),$$

and hence $\text{var}(g) \leq C u |v| / \log(1/u)$. Therefore, there is a constant C' such that

$$\mathbf{P}\left[f(x) > s(u) + C' \sqrt{|v| / \log(1/u)}\right] < u/2.$$

That is, $s(u/2) \leq s(u) + C' \sqrt{|v| / \log(1/u)}$. Induction gives for $n = 1, 2, \dots$

$$s(2^{-n}) \leq s(1/2) + O(1) \sqrt{n |v|},$$

which is the “upper tail” bound from (1). The lower tail is treated similarly. (This proof is fairly general. Using the more specific arguments from Section 3, a slight improvement for the tail estimates in certain ranges may be obtained.)

Acknowledgements. We are most grateful to Elchanan Mossel for very useful discussions and to Jan Vondrák for detecting a mistake in what was Theorem 2 in a previous version of the paper.

References

- [1] J. Baik, P. Deift and K. Johansson, On the distribution of the length of the longest increasing subsequence of random permutations, *J. Amer. Math. Soc.* **12** (1999), 1119–1178.
- [2] W. Beckner, Inequalities in Fourier analysis, *Annals of Math.* **102** (1975), 159–182.
- [3] S. G. Bobkov and C. Houdre, A converse Gaussian Poincare-type inequality for convex functions, *Statist. Probab. Lett.* **44** (1999), 281–290.
- [4] A. Bonami, Etude des coefficients Fourier des fonctions de $L^p(G)$, *Ann. Inst. Fourier* **20** (1970), 335–402.
- [5] J. Bourgain, J. Kahn, G. Kalai, Y. Katznelson and N. Linial, The influence of variables in product spaces, *Isr. J. Math.* **77** (1992), 55–64.
- [6] J.-D. Deuschel, O. Zeitouni, On increasing subsequences of I.I.D. samples, *Combin. Probab. Comput.* **8** (1999), 247–263.
- [7] R. Durrett, Harry Kesten’s publications: a personal perspective, in *Perplexing problems in probability*, Progr. Probab., 44, Birkhäuser Boston, Boston, MA, 1999, 1–33.
- [8] E. Friedgut, Influences in Product Spaces, KKL and BKKKL Revisited. Preprint.
- [9] K. Johansson, Transversal fluctuations for increasing subsequences on the plane, *Probab. Theory Related Fields* **116** (2000), 445–456.
- [10] J. Kahn, G. Kalai and N. Linial, The influence of variables on boolean functions, *Proc. 29-th Ann. Symp. on Foundations of Comp. Sci.*, (1988), 68–80.
- [11] H. Kesten, Aspects of first passage percolation, in *École d’été de probabilités de Saint-Flour, XIV—1984*, 125–264, Lecture Notes in Math., 1180, Springer, Berlin, 1986.
- [12] H. Kesten, On the speed of convergence in first passage percolation, *Ann. Applied Prob.* (1993), 296–338.

- [13] M. Ledoux, *The concentration of measure phenomenon*, Mathematical Surveys and Monographs, 89, American Mathematical Society, Providence, RI, 2001.
- [14] C. M. Newman and M. Piza, Divergence of shape fluctuations in two dimensions. *Ann. Probab.* **23** (1995), 977–1005.
- [15] R. Pemantle and Y. Peres, Planar first-passage percolation times are not tight in *Probability and phase transition (Cambridge, 1993)*, 261–264, Kluwer Acad. Publ., Dordrecht, 1994.
- [16] M. Talagrand, On Russo’s approximate zero-one law, *Ann. of Probab.* **22** (1994), 1576–1587.
- [17] M. Talagrand, Concentration of measure and isoperimetric inequalities in product spaces, *Inst. Hautes Études Sci. Publ. Math.* **81** (1995), 73–205.

THE WEIZMANN INSTITUTE OF SCIENCE, REHOVOT 76100, ISRAEL
 itai@wisdom.weizmann.ac.il
<http://www.wisdom.weizmann.ac.il/~itai/>

THE HEBREW UNIVERSITY, GIVAT RAM, JERUSALEM 91904, ISRAEL
 kalai@math.huji.ac.il
<http://www.ma.huji.ac.il/~kalai/>

MICROSOFT RESEARCH, ONE MICROSOFT WAY, REDMOND, WA 98052, USA
 schramm@microsoft.com
<http://research.microsoft.com/~schramm/>